


Health and Well-being of Family Carers in Ireland:

Results of a survey of recipients of the Carer's Allowance

Combat Poverty Agency Offices, Tuesday October 28th 2008
17:30-7pm.

Liam O'Sullivan, National Development Officer, Care Alliance Ireland
B. Social Studies, NQSW,
M.A. Youth and Community Studies, P.G. Cert. Social Research Skills.

Ann Stokes, Research and Policy Officer, Care Alliance Ireland
B Soc Sc, MA Soc Sc (Social Policy)



Promoting understanding. Informing policy.

Care Alliance Ireland

The National Network of Voluntary Organisations for Family Carers



- 66 Member Organisations – 11 Board Members
- We aim to support organisations in their direct work of supporting family carers all over Ireland
- We provide info, education & training regarding the needs of family carers
- We carry out research relating to family carers and promote inter agency collaboration

2

Background to Research

The context

- Funding – C.P.A, DSFA, CAI
- Timeframe – 1year – 3 years

3

This presentation will:

- Summarise other research on Family Carers Health
- Outline objectives of this piece of research
- Describe the Methodology used
- Present The Findings
- Ask 'Where do we go from here?'

4

The Hypothesis

- **Low Income full-time family carers are (hypothesised to be) a group that are particularly vulnerable to poor health**

5

Summary of Irish Research

- Limited, not recent
- Blackwell (1992) 1st quantifiable comparisons between carers and non-carers (29.5% carers vs 16.5% non carers at risk of depression)
- O'Connor & Ruddle (1988) caring and link to poor health
- SEHB (2000) caring negatively affects health of large proportion of carers
- O'Donoghue (2003) Importance of looking at health of care recipient and carer together
- Equality Authority (2004) recognition of health problems suffered by carers

6

Summary of UK Research

- More comprehensive
- Significantly greater levels of ill-health among carers than reported amongst the non-carer population, particularly for young carers (Carers UK, 2004)
- Positive correlation between intensity of caring and ill health (Maher & Green, 2002)
- Particularly high risk of depression and stress among carers (Boden, 2002; Carers UK, 2006)

7

Summary of International Research

- Higher rates of ill health among carers (O'Connell et al, 2003)
- Women experience more negative health effects (Morris, 2001)
- More inclined to worry or be stressed (Cummins, 2007)
- Carers less inclined to visit their GP (Evercare, 2006)

8

The Objectives

- Original scope:
 - to quantify reported family carer health as compared to the national non-carer population.

9

The Objectives (2)

- Scope evolved
 - Literature review
 - Mediators of carers health
 - Strategies used
 - Income
 - Household and Lifestyle Q's
 - General Profile

10

Sampling Method

- Population – 28,340
- Sample of 10% - 2,834
- Response rate 50% - sample of 1,413. (v's 11%ESRI,70%DSFA)
- Gender – 80% Female/20%Male
- A Comparative study (in part)
- Control for age, gender, educational attainment (proxy for class/income)

11

Socio-demographic profile of respondents

		% of carers
Age	16-24	21.1
	25-34	8.4
	35-44	23.5
	45-54	26.5
	55-64	22.5
	65+	15.1
Education	none/primary	27.2
	some secondary	37.7
	secondary	22.3
	some 3rd level	7.8
Marital status	completed 3rd level	4.8
	married/co-habiting	72.1
	widowed	2.8
Location	divorced/separated	6.6
	never married	18.4
	total	36.7
Household composition	urban	33.3
	1 person	1.6
	2	31.2
	3	23.5
	4	16.8
	5	13.0
	6	6.3
7+ persons	5.1	
Household with person <16 yrs		37.2

12

Income-related characteristics of the carers-

	% of carers	
Carer's Allowance	In receipt of full CA	86.1
	In receipt of partial CA	13.9
Medical card	Have a medical card	75.5
Other sources of income	No other sources of income	74.6
	Part-time job	9.6
	Odd job / occasional labour	1.8
	Other social welfare payments	3.3
	Income from relatives / friends	4.8
	Other	5.9
	Other	5.9
Spouse / partner status	Not living with spouse / partner	28.9
	Spouse full-time employed	26.4
	Spouse part-time employed	4.7
	Spouse unemployed	6.3
	Spouse receiving disability pay	17.0
	Spouse a pensioner	15.0
	Spouse in education/training	0.7
	Other	3.2


13

Amount and location of caring- mostly 1 caree >59hrs

	% of carers	
Number of people cared for	1	86.5
	2	12.9
	3	0.5
	4	0.1
Hours of care provided	0-19	1.7
	20-39	8.7
	40-59	13.6
	More than 59	76.0
Living with person cared-for	Co-resident	86.8
	Live in different households	13.2

14

Age of person being cared for- two peaks – children and over 70's



15

Main types of relationship with person cared for – 98.3% Family

	% of carers
Child	33.5
Child < 18	(18.6)
Child ≥ 18	(14.9)
Spouse/partner	25.6
Parent/parent-in-law	30.0
Other relative	9.3
Neighbour / friend / other	1.7

16

Self-reported general health ?



17

Self-reported general health

Health rating	Carers	SLÁN (weighted)
	per cent of carers	per cent of adult population
5 Excellent	10.5	12.9
4 Very good	25.5	32.0
3 Good	46.1	37.5
2 Fair	16.1	14.8
1 Poor	1.8	2.8
Total	100 per cent	100 per cent

(Base: total sample of carers, missing data = 13; χ^2 df, 4=46.6, p<.0001)

18

Satisfaction with health



19

Satisfaction with health

Satisfaction rating	Carers	SLÁN (weighted)
	per cent of carers	per cent of adult population
Very dissatisfied	3.4	2.5
Dissatisfied	9.3	9.2
Neither satisfied nor dissatisfied	17.1	18.2
Satisfied	55.4	53.1
Very satisfied	14.7	17.0
Total	100 per cent	100 per cent

(Base: total sample of carers, missing data = 40; $\chi^2_{df,4}=8.9$, n.s.)

20

Quality of life



21

Quality of life

QoL rating	Carers	SLÁN (weighted)
	per cent of carers	per cent of adult population
Very poor	2.0	1.2
Poor	6.5	2.8
Neither good nor poor	23.3	13.5
Good	52.1	54.8
Very good	16.0	27.7
Total	100 per cent	100 per cent

(Base: total sample of carers, missing data = 30; $\chi^2_{df,4}=177.6$, $p<.0001$)

22

Specific health problems experienced in past 12 months

Specific health problem	% of carers
Stress / nervous tension	40.8
Headaches	29.8
Lower back pain	26.3
Aching joints	25.8
Anxiety	22.6
Depression	17.6
High blood pressure	15.8
Urinary problems	8.4
Rheumatoid arthritis	8.2
Osteo-arthrits	7.7
Asthma	6.8
Diabetes	3.8
Chest/lung disease	2.6
Angina	1.8
Stroke	0.6
Cancer	0.6
Heart attack	0.4

(Base: total sample of carers for each specific health problem)

23

Impact of Caring On Health

Health has suffered	per cent of carers
Yes	29.5
No	70.5
Total	100 per cent

(Base: total sample of carers; missing data = 30)

- Ruddle and O'Connor - 30% believed that their health had suffered due to the demands of caring (1988)

24

Perceived impact of caring role on leisure/recreational activities



25

Perceived impact of caring role on leisure/recreational activities

Leisure limited	% of carers
Not at all	16.0
A little	38.6
Quite a lot	26.9
A great deal	18.5
Total	100 per cent

(Base: total sample of carers; missing data = 39)

26

Specific difficulties experienced by carers

Most difficult things about caring	% of carers
Sadness for person I care for	55.6
Being constantly on call	50.9
Stress	49.2
Emotional strain	43.1
Lack of sleep / tiredness	42.1
Frustration	31.7
Isolation	22.8
Guilt	14.7

(Base: total sample of carers, multiple response; missing data = 80)

27

Strategies to cope with caring role



28

Strategies to cope with caring role

Strategies used to keep going	% of carers
Talk to friends	65.7
Watch TV	62.5
Praying / Faith	39.6
Exercise	35.9
Smoking	19.0
Use respite	13.2
Take medication	11.1
Drink alcohol	9.0
Attend support group	6.8
Use phone line support	4.5
Alternative medicine	4.0

(Base: total sample of carers, multiple responses; missing data = 50)

29

Rural/Urban Differences



30

Rural/Urban Differences

- Rural >
 - Phone line support (4.7% v 3.9%)
 - Exercise (35.1% v 34.2%)
- Urban >
 - Prayer (40% v 36.4%)
 - Use alternative medication (4.1% v 3.6%)
 - Attend a support group (7.8% v 6%)
 - Use respite (13.6% v 11.8%)
 - Use medication (11.4% v 9.7%)
 - Smoke (21.7% v 15.8%)
 - Drink alcohol (10.2% v 7.8%)
 - Watch TV (62% v 59.7%)

31

Support

Support from household, wider family, friends and people in the workplace

Level of support	% of carers
Little/None	18.7
Some, but no major source	29.0
One major source	32.0
More than one major source	20.3
Total	100 per cent

(Base: total sample of carers)

32

Overall Comparative Support Score

Overall Comparative Support Score:

Survey	Mean	N	St Deviation
Carers	8.776	1413	5.98409
Slán (Weighted)	12.8647	5525	9.06480

33

Lifestyle - Smoking



34

Lifestyle - Smoking

Smoking cigarettes / cigars now	Carers	SLÁN (weighted)
	% of carers	% of adult population
Smoke regularly	25.9	21.0
Smoke occasionally	4.8	3.3
Do not smoke	69.3	75.7
Total	100.0	100.0

(Base: total sample of carers, missing data = 23; $\chi^2_{df,2}=24.4, p<.001$)

35

Lifestyle – Alcohol Consumption



36

Lifestyle – Alcohol Consumption

Last time had alcoholic drink	Carers	SLÁN (weighted)
	% of carers	% of adult population
During last week	30.1	54.6
One week to 1 month ago	18.6	13.8
One month to 3 months ago	9.4	5.6
Three months to 12 months ago	8.5	3.6
More than 12 months ago	11.5	5.9
Never had alcohol beyond sips or tastes	21.9	16.6
Total	100.0	100.0

(Base: total sample of carers, missing data = 53; $\chi^2df,5=296.6, p<.00001$)

37

Lifestyle – Alcohol Consumption (2)

Days drinking per week	Carers	SLÁN (weighted)
	% of carers	% of adult population
1	53.5	28.2
2	23.7	28.5
3	11.4	18.0
4	3.8	10.8
5	2.7	6.3
6	0.5	1.8
7	4.3	6.4
Total	100.0	100.0

(Base: people who drank in last 12 months; $\chi^2df,6=140.6, p<.00001$)

38

Injuries

Table 6.4. Injuries interfering with daily activities

One or more injuries in last 2 years serious enough to interfere with daily activities	Carers	SLÁN (weighted)
	per cent of carers	per cent of adult population
Any injury (Of which back injury)	14.0 (61.1)	13.0 ??
No injury	86.0	87.0
Total	100.0	100.0

(Base: total sample of carers, missing data = 55; $\chi^2df,1=2.0, n.s.$)

39


Factors associated with negative outcomes for carers

Factors increasing the likelihood of carer's reporting their health suffering due to being a carer

Dimension	Specific factor	Relative likelihood of reporting health issue suffered (compared with the first category in each case)
Type of care-related impact	1. Leisure limited by caring	
	- not at all	1
	- a little	1.8
	- quite a lot	4.9
	- a great deal	7.8
	2. Stress	
	- no	1
- yes	3.2	
3. Lack of sleep / tiredness	- no	1
	- yes	2.7
	4. Emotional strain	
- no	1	
- yes	2.7	
7. Isolation	- no	1
	- yes	1.7
Moderators	6. Talk to friends (as coping strategy)	
	- yes	1
- no	1.7	
Health-related lifestyle	5. Injury in last two years	
	- no	1
- yes	3.1	

40

Combined effects on health of key factors




41

Quality of Life – Stepwise Logistic Regression

- These variables, in order of statistical significance, were:
 - leisure limited by caring (great deal 3.3, quite a bit 3.0)
 - availability of support (little or none 2.7)
 - Stress (1.9)
 - talking to friends (not reported as coping strategy) (2.0)
 - Isolation (1.9)
 - spouse in not full-time employment (1.9)
 - lack of sleep / tiredness (1.7)
 - emotional strain (1.5)
 - Gender (1.6)
 - prayer / faith (as coping strategy) (1.5)
 - injury in the last two years (other than back injury). (2.1)


42

Additive effect of interference with leisure, stress and lack of support on quality of life among carers



43

Additive effects of hours of care, being constantly on call and type of needs on limitation of leisure/recreation activity



44

Summary - 1

- Less like to report Very Good or Excellent Health
- Lower Quality Of Life
- Similar Satisfaction with Health
- High levels of Stress, Headaches, Lower back Pain, Anxiety and Depression
- 30% report caring impacting on their health
- Difficulties include sadness for caree, constantly on call, tiredness

45

Summary - 2

- Coping strategies – Friends, TV, Prayer, Exercise
- Support levels – Comparatively low
- Higher levels of smoking
- Lower levels of alcohol consumption

46

Key Recommendations

1. Promoting awareness of Family Carers and their support needs
2. Reducing carer stress
3. Targetting Male Carers
4. Carer entitlements and Income
5. Promoting Carers Health


47

Where do we go from here?

- Third level Liasion – NUI Galway,UCC,TCD,WIT
- Family Carers Research Alliance
- Further Research
 - Further analysis of existing data – UCD ISSDA
 - SLAN 06 Data Qualitative Research
 - Qualitative – Mediating Factors,Meaning of Support, Expectations, use of Respite care
 - Male Carers
 - Testing of Hypothesis
 - Carers Association Research

48

Proposed Hypothesis Model of Causal Connections



49

Acknowledgements

- DSFA, CPA, Anglo Irish Bank
- Peers/Colleagues
- Family Carers

50

Contact Details

- Liam O'Sullivan – ndo@carealliance.ie
- Ann Stokes – research@carealliance.ie
- www.carealliance.ie
- + 353 1 874 7776



Final Report will be available on our website in coming weeks

51

Appendix 1 - Bibliography

- Blackwell, J., O'Shea, E., Moane, G. and Murray, P. (1992). *Care Provision and Cost Measurement: dependent elderly people at home and in geriatric hospitals*. Dublin: Economic and Social Research Institute.
- Boden C. (2002) *CASS Carers Survey, Depression and Carers*. Carers Association Southern Staffordshire.
- Care Alliance Ireland (2008, unpublished) *Health and Well-being of Family Carers in Ireland: Results of a survey of recipients of the Carer's Allowance*.
- Carers UK (2004) *In Poor Health: The impact of caring on health*. Carers UK Research Report.
- Cullen et al (2004) *Caring Work and Public Policy*. Work Research Centre. Dublin: The Equality Authority.
- Cummins, R (2007). Article in *The Age*, October 15, 2007 <http://www.theage.com.au/news/national/warning-signs-for-carers-wellbeing/2007/10/15/1192320891253.html> Website accessed April 10th 2008.
- Evercare (2006) (in collaboration with National Alliance for Caregiving) *Study of Caregivers in Decline A Close-up Look at the Health Risks of Caring for a Loved One* Findings from a National Survey
- Kelleher, C, Nic Gabhainn, S., Friel, S., Corrigan, H., Nolan, G., Sixsmith, J., Walsh O., Cooke, M. (2003) *Slán, Results of National Health and Lifestyle Surveys SLAN (Survey of Lifestyle, Attitudes and Nutrition)* Centre for Health Promotion Studies, Health Promotion Unit, Department of Health and Children, Dublin.
- Maher J and Green, H (2002). *Carers 2000*. Office for National Statistics, The Stationery Office.

52

Bibliography (2)

- Morris, M. (2001) *Gender-sensitive home and community care and caregiving research: a synthesis paper*, Ottawa: Health Canada Women's Health Bureau.
- O'Connell, B., Bailey, S. and Walker A. (2003) *Promoting the health and well being of older carers: A proactive strategy* Aust Health Rev 2003; 26 (2): 7866
- O'Connor, J and Ruddle, H (1988a) *Caring for the Elderly Part 1- A study of carers at home and in the community. Report No. 18*, Dublin: National Council for the Aged
- O'Connor, J and Ruddle, H (1988b) *Caring for the Elderly Part II – The Caring Process: a study of carers in the home. Report No. 19*, Dublin: National Council for the Aged
- O'Donoghue, (2003) (Unpublished) *Research project on Carers*. Supported by Eiri Corca Baiscinn.
- South Eastern Health Board (2000). *Listening to the Voices of Carers: an exploration of the health and social care needs and experiences of informal carers of older people*. Kilkenny: South Eastern Health Board.

53